Declassified in Part - Sanitized Copy Approved for Release @ 50-Yr 2013/07/23 : CIA-RDP83-00423R000700410007-2

20862

50X1

INTRA-OFFICE MEMORANDUM

TO: [redacted]

DATE: March 3, 1953

FROM:

SUBJECT: New Products - February, 1953

Domestic

Aureomycin Packing and Dressing

Description: These two products are being presented for local antibacterial therapy. According to the maker, Aureomycin dressing is non-adherent, non-macerating with wide spectrum antibacterial action; and the packing is suggested for use wherever plain or impregnated gauze packing was formerly used to control infection for hemostatis, drains etc.

50X1

Source: [redacted]

Distrycin (Also introduced in Lima, Peru under the name **Ambistryn**)

Description: Each vial contains:
Streptomycin Sulfate and
Dihydrostreptomycin Sulfate in
equal parts.

Indications: The same as streptomycin and dihydrostreptomycin but less toxic.

50X1

Form Supplied: [redacted]

Source: [redacted]

Fumidil

Description: Each capsule contains:
Fumagillin 10 mg.

Indications: For treatment of amebic or tropical dysentery. It kills Endamoeba histolytica directly. It has no antibacterial spectrum and does not disturb the normal flora of the gastro-intestinal tract.

50X1

Form Supplied: Bottles of 20, 30 and 60.

Source: [redacted]

Remanden

Description: Each tablet contains:
Potassium Penicillin G. Crystalline 100,000 u.
Probenecid 0.25 gm.

Declassified in Part - Sanitized Copy Approved for Release @ 50-Yr 2013/07/23 : CIA-RDP83-00423R000700410007-2

Declassified in Part - Sanitized Copy Approved for Release @ 50-Yr 2013/07/23 : CIA-RDP83-00423R000700410007-2

Indications:	For use when prolonged plasma concentrations of penicillin are needed. It acts to prolong blood levels of penicillin and allows for an increase of penicillin plasma concentrations two to four times as compared to conventional oral doses.
Form Supplied:	Vials of 12.
Source:	

50X1

Declassified in Part - Sanitized Copy Approved for Release @ 50-Yr 2013/07/23 : CIA-RDP83-00423R000700410007-2

Declassified in Part - Sanitized Copy Approved for Release @ 50-Yr 2013/07/23 : CIA-RDP83-00423R000700410007-2

Foreign

Antibiotic Powder (Sterile)

Description: Each sterile envelope (1 Gm.) contains:

Potassium Penicillin G	10,000 I.U.
Dihydrostreptomycin	20,000 mcgm.

Indications: Abdominal and other types of general surgery and following debridement of wounds. Saprophytic ulceration associated with carcinoma, or following necrosis of tissues from radium radio-therapy or electrocautery. Purulent infections of the skin and superficial wounds (chronic vaginitis, secondary infections accompanying "Trichomonas vaginalis and Monilia infections). Pre- and post-operatively in the genital tract.

50X1

Form Supplied: Boxes of 6, 25 and 100 sterile envelopes.

Source: [redacted].

Bismucilna

Description: Each 3 cc. vial contains:

Procaine Penicillin	300,000 U.
Bismuth	0.09 Gm.

in an oil suspension with AMS

Indications: Syphilis, amygdalitis and acute pharyngitis.

Administration: 1 dose every four days or 1 dose every two days in certain cases of syphilis

Form Supplied: Packages of 1 or 6 dose vials.

Source: [redacted] 50X1

Isacilin-S

Description: Each dose contains:

Procaine Penicillin G	300,000 U.O.
Potassium Penicillin G, crystalline	100,000 U.O.
Dihydrostreptomycin Sulfate	.5 Gm.

Indications: Meningitis and pneumonia caused by Hemophilus influenzae, Escherichia coli and Klebsiella pneumoniae.
Acute osteomyelitis, abscesses of soft tissues, thrombosis of sinus cavity, empyema, and kidney abscesses.
Gaseous gangrene, malignant edema
Cellulitis, mastoiditis, puerperal infection, peritonitis.
Blennorrhea, arthritis, endocarditis, prostatitis, epididymitis.

Source: [redacted]

50X1

Declassified in Part - Sanitized Copy Approved for Release @ 50-Yr 2013/07/23 : CIA-RDP83-00423R000700410007-2

Declassified in Part - Sanitized Copy Approved for Release @ 50-Yr 2013/07/23 : CIA-RDP83-00423R000700410007-2

Micocillina
Description: Selected strains of Penicillin Notatum
Indications: Superficial and profound lesions, wounds, abscesses, sycosis, empyema, purulent otitis, ulcers.
Form Supplied: Liquid - Vial of 2, 5 and 10 cc.
Powder Tube of 5 gr.
Ointment Tube of 10 gm.
Vaginal Suppositories - Boxes of 10
Tablets Tube of 20 (Trade name "Mico").
50X1
Source:

Penisulmida
Description: Contains:
Penicillin
Sulfadiazine
Sulfamerazine
Sulfamethazine
Indications: Augments the efficiency of the sulfonamides, diminishes the danger of crystallura and lessens the frequency of allergies.
50X1
Source:

Pen-Valeas
Description: Each suppository contains:
Penicillin
Bismuth
Cobalt
Indications: Used in treatment of the following infections:
Spirillum, micrococcus, streptococcus, gonococcus, pneumococcus, staphylococcus.
Source: 50X1

PGA 500
Description: Each tablet contains:
Penicillin G with 500,000 U.
ammonium
When a sufficient quantity of this penicillin is administered orally, the blood concentration is elevated rapidly and attains a maximum within 1 hour.
50X1
This new therapy maintains the blood level for about six hours.
Form Supplied: Bottles of 12 and 100 tablets.
Source:

Seclomycin
Description: Each dose contains sodium and procaine penicillin plus streptomycin sulfate.
Indications: Effective against the "mixed" infections commonly occuring in peritonitis, infected wounds, urinary tract etc.

Declassified in Part - Sanitized Copy Approved for Release @ 50-Yr 2013/07/23 : CIA-RDP83-00423R000700410007-2

Declassified in Part - Sanitized Copy Approved for Release @ 50-Yr 2013/07/23 : CIA-RDP83-00423R000700410007-2

Form Supplied: Dry form for aqueous injections and in single-dose vials.

Source: 50X1 [redacted]

SP3 Sulfapenisil

Description: Each tablet contains

Sulfadiazine	0.165 gr.
Sulfamerazine	0.165 gr.
Sulfamethazine	0.165 gr.
Crystalline Penicillin	50,000 u.

50X1

Source: [redacted]

Streptocol

Description: Each 5 gr. tablet contains:
Streptomycin
3 - Sulfonamides
Vitamin PP
Vitamin K

Indications: Enteritis and acute enterocolitis, subacute and chronic coli bacteria infections, dysentery, salmonella, infantile diarrhea, tuberculous diarrhea etc. For pre- and post-operative use in intestinal surgery.

Administration: For enteritis and enterocolitis

Adult 2 to 4 tablets at one time followed by 1 to 2 tablets every five hours.

Children from one to three years old, half dose.

For pre- and post-operative use:

Adult 2 tablets every two hours for five times (in particular cases for 10 times) beginning the morning before the operation.

50X1

Children from one to three years old, half dose.

Form Supplied: Bottles of 10 tablets

Source: [redacted] ltico

Tampovagan Pessaries with Penicillin (5,000 i.u.)

Description: Each globule contains 5,000 i.u. of penicillin in a cocoa-butter base.

Indications: All cases of vaginal fluor in which the causative organism is known to be

Declassified in Part - Sanitized Copy Approved for Release @ 50-Yr 2013/07/23 : CIA-RDP83-00423R000700410007-2

Declassified in Part - Sanitized Copy Approved for Release @ 50-Yr 2013/07/23 : CIA-RDP83-00423R000700410007-2

sensitive to penicillin, in cases showing an idiosyncrasy toward sulfa drugs, or where sulfonamides have been previously employed unsuccessfully.

Tampovagan P.S.S.

Description: These contain 5,000 u. of penicillin with 0.25 g. of sulfanilamide and sulfathiazole.

Indications: All primary and secondary infections due to organisms sensitive to penicillin and/or sulfonamides. Gonorrhoea, syphilis, streptostaphylo and pneumococci, etc.

50X1

Form Supplied: Boxes of 8.

Source: [redacted]

For your information: The following products are not new, but they are of interest. Two Italian manufacturers of Chloramphenicol have combined it with sulfanilamide.

Sulfamicetina

Description: Each tablet contains:

Chloramphenicol	.1 g.
Sulfathiazole	.3 g.
Sulfapyridine	.1 g.

Indications: Antibacterial activity against a major number of germs, also effective in cases of resistance to other medicines.

Administration: 4 to 8 tablets daily in broken doses

Form Supplied: Package of 10 tablets

Source: [redacted]

50X1

Tetracetina

Description: Each tablet contains:

Tetraseptale (Polysulfamide)	.3 g.
Farmicetina (Chloramphenicol)	.1 g.

Indications: Sinus, Otomastoiditis, tonsillitis, oral infections, atypical pneumonia, broncho-pneumonia, angiocolitis, coleocysitits, entercolitis, dysentery, post-operatively in abdominal surgery, urinary infections etc.

Form Supplied: Bottles of 10 and 20 tablets

Source: [redacted]

50X1

Declassified in Part - Sanitized Copy Approved for Release @ 50-Yr 2013/07/23 : CIA-RDP83-00423R000700410007-2